

# *Thismia aliasii* (Thismiaceae), a new species from Terengganu, Peninsular Malaysia

Mat Yunoh Siti-Munirah<sup>1</sup>, Shakri Mohamad Alias<sup>2</sup>

<sup>1</sup> Forest Research Institute Malaysia, 52109 Kepong, Selangor, Malaysia

<sup>2</sup> Jabatan Perhutanan Negeri Terengganu, 21500 Kuala Terengganu, Terengganu, Malaysia

Corresponding author: Mat Yunoh Siti-Munirah (sitimunirah@frim.gov.my)

## Abstract

A new mycoheterotrophic species, *Thismia aliasii*, is described and illustrated. This species inhabits a hill dipterocarp forest in mountains of eastern Peninsular Malaysia. *Thismia aliasii* differs from other *Thismia* species by the following features: tepals equal in size and shape with different length of their appendages, appendages of the outer tepals shorter than those of the inner tepals (3.5 mm vs. ca. 26–32 mm long), stamen supraconnective at apex with three long filiform appendages and two acute appendages, and the margins of individual connectives abaxially raised into a conspicuous rib. With respect to floral morphology, *T. aliasii* should be placed to *Thismia* subsect. *Odoardoa*. According to the categories and criteria of the IUCN Red List, *T. aliasii* is provisionally classified as Critically Endangered (CR).

**Key words:** Achlorophyllous plants, endemism, Gunung Chemerong, monocots, Terengganu, *Thismia* subsection *Odoardoa*



Academic editor: Michal Hroneš

Received: 1 September 2024

Accepted: 20 February 2025

Published: 31 March 2025

**Citation:** Siti-Munirah MY, Mohamad Alias S (2025) *Thismia aliasii* (Thismiaceae), a new species from Terengganu, Peninsular Malaysia. PhytoKeys 254: 175–188. <https://doi.org/10.3897/phytokeys.254.136085>

Copyright:

© Mat Yunoh Siti-Munirah & Shakri Mohamad Alias. This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

## Introduction

The fully mycoheterotrophic monocot family Thismiaceae is mainly confined to tropical and subtropical regions. Species of the family are small herbs that usually inhabit shady, humid environments, often hidden under the leaf litter (Merckx et al. 2024). The genus *Thismia* Griff., often referred to as "fairy lanterns" (Wapstra et al. 2005), is remarkable due to its unusual flower appearance. The unique flower design of the genus is related to the specialized pollination mechanisms that involve small insects such as fungus gnats (belonging to the genus *Corynoptera*) (Guo et al. 2019). *Thismia* flowers were also visited and possibly pollinated by a wide range of other invertebrates, such as scuttle flies (Phoridae, Diptera) (Yudina et al. 2021).

The genus *Thismia* comprises about 116 species (Besi et al. 2024; Chung et al. 2024; Chunyang et al. 2024; Nuraliev and Sennikov 2024; POWO 2024; Siti-Munirah et al. 2024; Ya et al. 2024). It is the most widespread and species-rich genus within the family and has a high degree of endemism. Its range extends from tropical and subtropical Asia to northern and eastern Australia and New Zealand, and from Costa Rica to tropical South America with an isolated occurrence in North America (POWO 2024). Many of its species are considered extremely rare, with scattered

distributions, and some may already be extinct, but there are also species with quite wide ranges and multiple known localities (Dančák et al. 2020).

The state of Terengganu is currently known to be the richest of the Peninsular Malaysian states in the species diversity of *Thismia* (Table 1, Fig. 1). All the 13 species known from Terengganu are present in the northern part of the state. The southern Terengganu, in contrast, is known to be inhabited by only two species of *Thismia*, both found in the vicinity of hill and mountain range known as Gunung (Gn.) Chemerong, which is part of the Hutan Lipur Gn. Chemerong (HLGC) or Chemerong Forest Eco Park and also well known as Chemerong-Berembun-Langsir (CBL) mountain range. This area is, in turn, a part of the Pasir Raja Forest Reserve (FR). One of these two species is *T. aseroe* (Fig. 2D), whereas the other one is *T. aliasii* described here as new to science.

Materials and methods

The assessment is based on the material collected in Gn. Chemerong, Hulu Terengganu, Terengganu (Fig. 3). *Thismia aliasii* was first discovered during a field trip of the second author to CBL in 2019. The specimens were preserved in 70% ethanol and stored in the Kepong Herbarium (KEP). Three flowering individuals and one fruiting plant were collected and studied in total. Morphological characteristics were examined and measurements performed using an Olympus SZ61 stereomicroscope and high-resolution macrophotography. The measurements were made on fresh and spirit material.


Taxonomic account

*Thismia aliasii* Siti-Munirah, sp. nov.  
urn:lsid:ipni.org:names:77359488-1  
Figs 4–6

**Diagnosis.** *Thismia aliasii* is very similar to the species of the *T.* subsection *Odoardoa*, as the tepals are the same in shape and size. However, the tepal appendages of the new species are of unequal length, the inner ones are longer than the outer ones, while the tepal appendages of the other species are of


Table 1. *Thismia* of Terengganu state (Siti-Munirah and Dome 2019, 2021, 2022, 2023a, 2023b; Siti-Munirah et al. 2021).

No.	Species	Locality in Terengganu	Status of Endemism
1.	<i>T. alba</i> Holttum ex Jonker	Taman Negeri Kenyir, Tembat FR	No
2.	<i>T. aliasii</i> Siti-Munirah	Gn. Chemerong (Pasir Raja FR)	Endemic
3.	<i>T. arachnites</i> Ridl.	Taman Negeri Kenyir	No
4.	<i>T. aseroe</i> Becc.	Gn. Chemerong (Pasir Raja FR)	No
5.	<i>T. brunneomitroides</i> Suetsugu & Tsukaya	Taman Negeri Kenyir	No
6.	<i>T. clavigeroides</i> Chantanaorr. & Seelanan	Taman Negeri Kenyir	No
7.	<i>T. domei</i> Siti-Munirah	Hulu Telemong FR	Endemic
8.	<i>T. javanica</i> J.J.Sm.	Hulu Telemong FR, Tembat FR	No
9.	<i>T. kenyirensis</i> Siti-Munirah & Dome	Taman Negeri Kenyir	Endemic
10.	<i>T. latiffiana</i> Siti-Munirah & Dome	Gn. Sarut (Hulu Nerus FR)	Endemic
11.	<i>T. ornata</i> Dančák, Hroneš & Sochor	Gn. Padang	No
12.	<i>T. sitimeriamiae</i> Siti-Munirah, Dome & Thorogood	Gn. Sarut (Hulu Nerus FR)	Endemic
13.	<i>T. terengganuensis</i> Siti-Munirah	Hulu Telemong FR	Endemic



**Figure 1.** *Thismia* species occurring in the state of Terengganu **A** *Thismia alba* **B** *T. arachnites* **C** *T. aseroe* **D** *T. domei* **E** *T. javanica* **F** *T. kenyirensis* **G** *T. latiffiana* **H** *T. sitimeriamiae* **I** *T. terengganuensis*. Photos by Siti-Munirah (**A–G, I**) and Dome (**H**).






**Figure 2.** Landscapes of Gn. Chemerong and the species of *Thismia* found there **A** the Chemerong waterfall with a height of 370 m **B** the Chemerong River not far from the habitat of *T. aliasii* **C** *Thismia aliasii*, an individual discovered by Mohamad Alias in 2019 (FRI 91119) **D** *Thismia aseroe* (FRI 79116) found on the trail at Hutan Lipur Chemerong. Photos by Siti-Munirah (**A, B, D**) and Mohamad Alias (**C**).


equal length. In addition, in the new species the margins of the individual connectives are raised abaxially into the conspicuous rib, whereas connectives are almost flat abaxially in the rest of the species.

**Type.** MALAYSIA. • Peninsular Malaysia: Terengganu, Hulu Terengganu District, Hutan Simpan Pasir Raja, Chemerong Forest Eco Park, Gunung Chemerong, 4°39'33.2"N, 102°58'58.6"E, elev. ca 640 m, 26 July 2023, Siti-Munirah, FRI 79119 (holotype KEP!, spirit collection, barcode no. SC13201).



**Figure 3.** Gunung Chemerong (black circle) in Terengganu, the only known locality of *Thismia aliasii*.


**Description.** Achlorophyllous herb, up to ca. 11 cm tall, mostly glabrous (where not stated otherwise). **Roots** vermiform, unbranched, ca. 1.5 mm in diameter, light brown. **Stem** erect, up to 56 mm long, 1.8–2 mm in diameter, white-cream to light brown, bearing 1–2 flowers. **Leaves** up to 5, spirally arranged (arranged denser at stem base and looser at apex), triangular to narrowly triangular (shorter at stem base and longer and narrower at stem apex), scale-like, apex pointed, margin almost entire to slightly irregularly serrate, up to 7 mm long, ca. 2 mm wide at base, colored similar to stem. **Flowers** terminal and solitary, or in 2-flowered terminal inflorescence, actinomorphic, ca. 58 mm long (including ovary, floral tube, tepals and tepal appendages when erected). **Involucral bracts** 3, similar to upper leaves, triangular to narrowly triangular, scale-like, acute, with entire margin, 8 mm long, ca. 2.5 mm wide at base, white-brownish/light brown. **Pedicel** to 1.5 mm long at anthesis, to ca. 4 mm long after anthesis, white-brownish. **Floral tube** obovoid funnel-shaped, 20 mm long, ca. 4 mm wide at base, ca. 8 mm wide at middle, ca. 10 mm wide distally; **outer surface** with minute glands, orange to sepia-brown, with 12 darker longitudinal ribs; **inner surface** smooth or rough, almost similar color to the outer



**Figure 4.** *Thismia aliasii* **A** plant with flower and roots **B** top view of flower showing tepals and annulus **C** bract (abaxial view) **D** leaf (abaxial view) **E** style and stigma **F** stamens (outer view) **G** stamens (inner view) **H** fruiting plant. All drawn by Mohamad Aidil Noordin from spirit material, FRI 91119.

surface, with transverse bars. **Tepals** 6, free, spreading, triangular, apex acute, 7 mm long, ca. 1–1.5 mm wide (ca. 1.5 mm at base), smooth, color similar to flower tube (dark brown at apex, yellow-orange at back), each apically bearing a tentacle-like appendage 0.5 mm wide and narrowing towards the apex; **appendage of outer tepal** up to 3.5 mm long, dark brown; **appendage of inner tepal** ca. 26–32 mm long, brownish. **Annulus** moderately raised, hexagonal in outline,





**Figure 5.** *Thismia aliasii* **A** flowering plant (over-brightened by a camera flash) **B** anthetic flower, top view **B1** annulus, top view **B2** tip of appendage of outer tepal **B3** tip of appendage of inner tepal **C** distal portion of flower, lateral view **C1** flower, top view **D** floral tube (with its apex removed), side view **D1** outer surface of floral tube (showing tiny glands) **E** inner surface of flower tube, pistil and ovary **E1** longitudinal section of ovary and pistil with stigma **F** fruiting plant **F1** seeds in capsule **F2** seeds. Photos by Siti-Munirah (**A**, **C1**: FRI 79119; **A1**, **B–E**: FRI 79167; **E1**, **F1**, **F2**: FRI 91119, spirit material) and Mohamad Alias (**F**: FRI 91119).

ca. 6 mm in diameter, with ring width ca 1.5 mm, brown-orange, aperture ca. 3 mm in diameter. **Stamens** 6, pendulous from annulus; **filaments** laterally pale orange, otherwise translucent white; **connectives** broad and flattened, fused laterally into a tube, shortly papillose, translucent white, with hairs around thecae, with margins abaxially raised to form conspicuous ribs along the sutures between the connectives, ribs distally slightly protruding beyond the apex of supraconnective; **interstaminal glands** conspicuous, placed between bases of lateral appendages; **supraconnective** with 3 filiform apical appendages (central appendage ca. 2 mm long and paired appendages ca. 2.4 mm long) and with 2 acute hook-like subapical appendages ca. 0.6 mm long positioned adaxially between the apical appendages and the lateral appendage; **lateral appendage** skirt-like, projecting towards the floral tube and not reaching the supraconnective apex, translucent white and brown on sides, lateral margin wavy. **Ovary** red, pale cream proximally and blackish distally; unilocular; **placentas** 3, free, column-like, arising at ovary base; ovules numerous. **Style** dark blackish-orange, ca. 1.2 mm long; stigma ca. 1.4 mm long, papillose, 3-lobed, erect, lobes  $\pm$  rectangular, bifurcate at the apex, black-greenish. **Fruit** dehiscent, cup-shaped, ca. 6 mm high, ca. 7 mm in diameter, pale brownish, darker in the upper part. **Seeds** long fusiform ca. 0.75 mm long, ca. 0.25 mm wide.

**Additional specimens examined.** MALAYSIA. • Peninsular Malaysia: Terengganu, Hulu Terengganu District, Pasir Raja FR, Chemerong Forest Eco Park, Gunung Chemerong, elev. ca 642 m, 3 October 2019, Mohamad Alias, FRI 91119 (KEP spirit collection, barcode no. SC13202); elev. ca 640 m, 26 July 2023, Siti-Munirah, FRI 79167 (KEP spirit collection, barcode no. SC13203).

**Distribution.** Endemic to Terengganu, Peninsular Malaysia. Currently only known from the type locality, Gunung Chemerong (Figs 2, 3, 7).


**Ecology.** The species inhabits moist shady areas of upper hill dipterocarp forest on moist soil at elevation of 640 m a.s.l. Flowering and fruiting recorded in July and October. The population was found in moist, shaded areas next to the main hiking trail to the summit of Gunung Chemerong Berembun Langsir at altitudes of 640 m above sea level. The species was found a few hundred meters away from the main river.

**Etymology.** The species is named after Mr. Alias (the second author), a current ranger of the Terengganu Forestry Department (JPNT), who is also a freelance photographer (known as John Sp) and who was the first to discover the species.

**Conservation status.** Since 2019, several surveys have been conducted at an area of 4 hectares, but the species was observed only twice, with a total of 5 individuals recorded. The main threat to the population is the degradation of habitat quality due to hiking activities, as the population is found near the main hiking trail which is heavily used by hikers ascending to the summits of Gunung Chemerong Berembun Langsir. Therefore, according to IUCN Red List (IUCN 2024) this species is assessed as Critically Endangered, CR B2ab(iii), D. More surveys are needed to determine the population size of the species.

**Notes.** *Thismia aliasii* is easily recognized within the genus by the combination of the following characters: vermiform roots, almost uniform flower coloration (light to dark orange to sepia-brownish red), inner tepals free from each other, unequal tepal appendages (appendages of the outer tepals being shorter than those of the inner tepals), stamens each with 3 long filiform apical appendages, 2 acute subapical appendages and a lateral appendage, and connectives





**Figure 6.** *Thismia aliasii* **A** longitudinally dissected stamen tube showing the margins of individual connectives abaxially raised into conspicuous ribs (spirit material) **A1** stamens, view from inside **B, C** stamens, view from outside **D, E** stamen tube, oblique view from below and view from below (showing apical appendages and longitudinal ribs). Yellow arrows: filiform apical appendages; red arrows: acute hook-like subapical appendage; orange arrows: anthers. All photos by Siti-Munirah (**A**: FRI 91119; **A1, B–E**: FRI 79167).



**Figure 7.** Habitat and habit of *Thismia aliasii* **A** plants in situ **B** plant with flower bud **C** plant with anthetic flower. All photos by Siti-Munirah.

laterally thickened into conspicuous abaxial interstaminal ribs. Within the infrageneric classification of Kumar et al. (2017), *T. aliasii* is assigned here to *Thismia* subgenus *Thismia* section *Thismia* subsection *Odoardoa* Schltr., as long as it has free tepals equal in shape and size. The new species probably



belongs to clade 5 defined by Shepeleva et al. (2020), which is characterized by free inner tepals and the presence of appendages of the outer and inner tepals. In addition, the fact that the appendages of the inner tepals are much longer than those of the outer tepals make *T. aliasii* similar to *T. neptunis* Becc., (Sochor et al. 2018), which belongs to *T.* subgen. *Thismia* sect. *Thismia* subsect. *Brunonithismia* Jonker according to Kumar et al. (2017), but was recovered as a member of clade 5 by Shepeleva et al. (2020).

The connectives of most *Thismia* species are usually flat and without any processes abaxially, making *T. aliasii* unique not only among *T.* subsect. *Odoardoa* but within the entire genus. The only other exceptions are species from *T.* sect. *Geomitra* (Becc.) Kumar & S.W. Gale which have ribs in the center of each stamen but differ by coralliform roots and flower mitre.

We summarize *Thismia* subsect. *Odoardoa* to comprise 23 species based on the above-mentioned characteristics and following the classifications of Kumar et al. (2017) and Shepeleva et al. (2020) with additions from Chantanaorrapint et al. (2016), Hroneš et al. (2018), Nishioka et al. (2018), Dančák et al. (2020), Siti-Munirah and Dome (2019), Siti-Munirah et al. (2024), Ya et al. (2024):

### **A checklist of *Thismia* subg. *Thismia* sect. *Thismia* subsect. *Odoardoa* Schltr., 1921.**

Type species: —*Thismia aseroe* Becc., 1878.

Species included:

- 1) *T. alba* Holttum ex Jonker, 1948.
- 2) *T. aliasii* Siti-Munirah, 2025. (this study)
- 3) *T. annamensis* K.Larsen & Aver., 2007.
- 4) *T. aseroe* Becc., 1878.
- 5) *T. bifida* M. Hotta, 1967.
- 6) *T. bryndonii* Tsukaya, Suetsugu & Suleiman, 2017.
- 7) *T. chrysops* Ridl., 1895.
- 8) *T. claviformis* Chantanaorr. & J.Wai, 2016.
- 9) *T. cornuta* Hroneš, Sochor & Dančák, 2018.
- 10) *T. domei* Siti-Munirah, 2019.
- 11) *T. filiformis* Chantanaorr., 2012.
- 12) *T. fumida* Ridl., 1890.
- 13) *T. grandiflora* Ridl., 1895.
- 14) *T. hexagona* Dančák, Hroneš, Kobrlová & Sochor, 2013.  
*T. hexagona* var. *grandiflora* Tsukaya, Suleiman & H.Okada, 2014.
- 15) *T. inconspicua* Sochor & Dančák, 2017.
- 16) *T. kinabaluensis* T.Nishioka & Suetsugu, 2018.
- 17) *T. lauriana* Jarvie, 1996.
- 18) *T. malayana* Siti-Munirah, Hardy-Adrian, Mohamad-Shafiq & Irwan-Syah, 2024.
- 19) *T. mullerensis* Tsukaya & H.Okada, 2005.
- 20) *T. ophiuris* Becc., 1878.
- 21) *T. ornata* Dančák, Hroneš & Sochor, 2020.
- 22) *T. pallida* Hroneš, Dančák & Rejzek, 2018.
- 23) *T. racemosa* Ridl., 1915.



Ten species of the subsection occur in Peninsular Malaysia (*T. alba*, *T. aseroe*, *T. chrysops*, *T. domei*, *T. fumida*, *T. grandiflora*, *T. malayana*, *T. ornata*, *T. racemosa* and *T. aliasii*), with five of them (*T. alba*, *T. aliasii*, *T. aseroe*, *T. domei* and *T. ornata*) occurring in the state of Terengganu. Whereas most of these species are characterized by tepal appendages equal in size and shape, in *T. aliasii* the appendages of the inner tepals are longer than those of the outer tepals. In addition, the stamen connectives laterally raised to form a conspicuous interstaminal rib are a unique feature of *T. aliasii* that is not found in the other species of *T.* subsect. *Odoardoa*.

## Acknowledgements

This research was conducted under the Flora of Peninsular Malaysia Project (FPM) at the Forest Research Institute Malaysia (FRIM), Kepong, funded by the Ministry of Natural Resources and Environmental Sustainability (NRES) under the 12<sup>th</sup> Malaysian Plan (SPPII No. P23085100018003) and the Nagao Research Grant to partially fund *Thismia* research. We thank the Jabatan Perhutanan Negeri Terengganu (JPNT) and Pejabat Hutan Daerah Dungun, MARDI team, as well as Dome Nikong of WildDome for their continuous support of *Thismia* research in Terengganu. Special thanks to Zubir Ahmad MI and Angan A for their help in searching for *Thismia* in the field and to the flora team (KEP (Imin K and Syahida-Emiza S) and KBG (Suhaida M, Khairil Azuar AK and Ahmad Norsidar AH)) and Zainul Ariffin AS and Amirul I for their support in the field work of Gn. Chemerong. We thank Michal Hroneš, Martin Dančák, Maxim Nuraliev and Sahut Chantanaorrapint for constructive comments on our manuscript, Mohamad Aidil Noordin for botanical drawings, Wendy SY Yong for assistance in conservation status assessment, and Mohd Faizal MT for encouragement and great help in many ways to the first author.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

This research was funded by the Ministry of Natural Resources and Environmental Sustainability (NRES) under the 12<sup>th</sup> Malaysian Plan (SPPII No. P23085100018003) and the Nagao Research Grant to partially fund *Thismia* research.

### Author contributions

All authors have contributed equally.

### Author ORCIDs

Mat Yunoh Siti-Munirah  <https://orcid.org/0000-0002-5062-9988>

### Data availability

All of the data that support the findings of this study are available in the main text.

## References

- Besi EE, Qistina Anis MJ, Syamil AR, Mohd Zulkifly S, Ag Shaffie AA, Go R (2024) *Thismia perlisensis* (Thismiaceae), a new red-annulus *Thismia* species from Peninsular Malaysia. *Phytotaxa* 675(1): 1–10. <https://doi.org/10.11646/phytotaxa.675.1.1>
- Chantanaorrapint S, Wai JS, Poopath M, Tetsana N (2016) *Thismia claviformis* (Thismiaceae), a new species from the Thai-Malay Peninsula. *Phytotaxa* 280(3): 292–296. <https://doi.org/10.11646/phytotaxa.280.3.8>
- Chung SW, Hsu TC, Lin CW, Yang TYA, Fanerii M, Pitisopa F, Li CW (2024) Three new species and new records of *Thismia* (Thismiaceae) in The Solomon Islands. *Taiwania* 69(3): 336–347. <https://doi.org/10.6165/tai.2024.69.336>
- Chunyang L, Haijun Y, Wenhao Q, Han X (2024) *Thismia jinzun* (Thismiaceae), A New Species of *Thismia* Genus from Hainan Island, China. *Phytotaxa* 664(4): 273–280. <https://doi.org/10.11646/phytotaxa.664.4.4>
- Dančák M, Hroneš M, Sochor M (2020) *Thismia ornata* and *T. coronata* (Thismiaceae), two new species from Sarawak, Borneo. *Willdenowia* 50(1): 65–76. <https://doi.org/10.3372/wi.50.50106>
- Guo X, Zhao Z, Shing Mar S, Zhang D, Saunders RMK (2019) A symbiotic balancing act: Arbuscular mycorrhizal specificity and specialist fungus gnat pollination in the mycoheterotrophic genus *Thismia* (Thismiaceae). *Annals of Botany* 124(2): 331–342. <https://doi.org/10.1093/aob/mcz087>
- Hroneš M, Rejžek M, Sochor M, Svátek M, Kvasnica J, Egertová Z, Pereira JT, Nilus R, Dančák M (2018) Two new species of *Thismia* subsect. *Odoardoa* (Thismiaceae) from Borneo. *Plant Ecology and Evolution* 151(1): 110–118. <https://doi.org/10.5091/plecevo.2018.1387>
- IUCN Standards and Petitions Committee (2024) Guidelines for Using the IUCN Red List Categories and Criteria. Version 16. Prepared by the Standards and Petitions Committee. <https://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Kumar P, Gale SW, Li J, Bouamanivong S, Fischer GA (2017) *Thismia nigricoronata*, a new species of Burmanniaceae (Thismieae, Dioscoreales) from Vang Vieng, Vientiane Province, Laos and a key to subgeneric classification. *Phytotaxa* 319(3): 225–240. <https://doi.org/10.11646/phytotaxa.319.3.2>
- Merckx VSFT, Gomes SIF, Wang D, Verbeek C, Jacquemyn H, Zahn FE, Gebauer G, Bidartondo MI (2024) Mycoheterotrophy in the wood-wide web. *Nature Plants* 10(5): 710–718. <https://doi.org/10.1038/s41477-024-01677-0>
- Nishioka T, Suetsugu K, Repin R, Kitayama K (2018) *Thismia kinabaluensis* (Thismiaceae), a new species from Mt. Kinabalu, Sabah, Borneo. *Phytotaxa* 360(2): 174–178. <https://doi.org/10.11646/phytotaxa.360.2.10>
- Nuraliev MS, Sennikov AN (2024) A checklist of *Thismia* section *Labiothismia* (Thismiaceae), reinstated to include Asian species earlier placed in the neotropical *T.* sect. *Glaziocharis*. *Phytotaxa* 662(1): 93–101. <https://doi.org/10.11646/phytotaxa.662.1.7>
- POWO (2024) Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. <https://powo.science.kew.org/> [Retrieved 1 September 2024]
- Shepeleva EA, Schelkunov MI, Hroneš M, Sochor M, Dančák M, Merckx VSFT, Kikuchi IABS, Chantanaorrapint S, Suetsugu K, Tsukaya H, Mar SS, Luu HT, Li H-Q, Logacheva MD, Nuraliev MS (2020) Phylogenetics of the mycoheterotrophic genus *Thismia* (Thismiaceae: Dioscoreales) with a focus on the Old World taxa: Delineation of novel natural groups and insights into the evolution of morphological traits. *Botanical Journal of the Linnean Society* 193(3): 287–315. <https://doi.org/10.1093/botlinnean/boaa017>

- Siti-Munirah MY, Dome N (2019) *Thismia domei* and *T. terengganuensis* (Thismiaceae), two new species, and *T. javanica*, a new record from Terengganu, Peninsular Malaysia. *PhytoKeys* 124: 123–137. <https://doi.org/10.3897/phytokeys.124.34728>
- Siti-Munirah MY, Dome N (2021) *Thismia* in Nerus Forest Reserve. *Conservation Malaysia Bulletin* 33: 8–9. <https://www.mybis.gov.my/pb/4663>
- Siti-Munirah MY, Dome N (2022) *Thismia latiffiana* (Thismiaceae), an unusual new species from Terengganu, Peninsular Malaysia. *PhytoKeys* 188: 105–114. <https://doi.org/10.3897/phytokeys.188.77061>
- Siti-Munirah MY, Dome N (2023a) *Thismia kenyirensis* (Thismiaceae), a new species from Taman Negeri Kenyir, Terengganu, Peninsular Malaysia. *PhytoKeys* 221: 61–72. <https://doi.org/10.3897/phytokeys.221.98571>
- Siti-Munirah MY, Dome N (2023b) *Thismia* (Thismiaceae) in Kenyir State Park, Terengganu. In: Ahmad et al. (Eds) *Proceeding of Kenyir State Park Scientific Expedition*. Universiti Malaysia Terengganu Press, Kuala Nerus, Terengganu, 49–56.
- Siti-Munirah MY, Dome N, Thorogood CJ (2021) *Thismia sitimeriamiae* (Thismiaceae), an extraordinary new species from Terengganu, Peninsular Malaysia. *PhytoKeys* 179: 75–89. <https://doi.org/10.3897/phytokeys.179.68300>
- Siti-Munirah MY, Hardy-Adrian C, Mohamad-Shafiq S, Irwan-Syah Z, Hamidi AH (2024) *Thismia malayana* (Thismiaceae), a new mycoheterotrophic species from Peninsular Malaysia. *PhytoKeys* 242: 229–239. <https://doi.org/10.3897/phytokeys.242.120967>
- Sochor M, Egertová Z, Hroneš M, Dančák M (2018) Rediscovery of *Thismia neptunis* (Thismiaceae) after 151 years. *Phytotaxa* 340(1): 71–78. <https://doi.org/10.11646/phytotaxa.340.1.5>
- Wapstra M, French B, Davies N, O'Reilly-Wapstra J, Peters D (2005) A bright light of the dark forest floor: observations of the fairy lanterns *Thismia rodwayi* F. Muell. (Burmanniaceae) in Tasmanian forests. University of Tasmania. Tasmanian Field Naturalist Club Inc 127: 2–18. <https://hdl.handle.net/102.100.100/590442>
- Ya J-D, Chen H-Y, Zhang W, Zhu R-B, Cai J, Yu W-B (2024) Phylogenetic and biogeographical analyses of *Thismia* (Thismiaceae) support *T. malipoensis* as the eighth species in China. –. *Willdenowia* 54(1): 47–63. <https://doi.org/10.3372/wi.54.54102>
- Yudina SV, Vislobokov NA, Nuraliev MS (2021) Evidences of a mixed pollination strategy in Vietnamese species of *Thismia* (Thismiaceae: Dioscoreales). *Wulfenia* 28: 109–128.